

Amendments to the Specification:

Please replace the paragraph at page 26, line 28 through page 27, line 26 with the following amended paragraph:

Other algorithms for determining homology or identity include, for example, in addition to a BLAST program (Basic Local Alignment Search Tool at the National Center for Biological Information), ALIGN, AMAS (Analysis of Multiply Aligned Sequences), AMPS (Protein Multiple Sequence Alignment), ASSET (Aligned Segment Statistical Evaluation Tool), BANDS, BESTSCOR, BIOSCAN (Biological Sequence Comparative Analysis Node), BLIMPS (BLOCKS IMPROVED Searcher), FASTA, Intervals & Points, BMB, CLUSTAL V, CLUSTAL W, CONSENSUS, LCONSENSUS, WCONSENSUS, Smith-Waterman algorithm, DARWIN, Las Vegas algorithm, FNAT (Forced Nucleotide Alignment Tool), Framealign, Framesearch, DYNAMIC, FILTER, FSAP (Fristensky Sequence Analysis Package), GAP (Global Alignment Program), GENAL, GIBBS, GenQuest, ISSC (Sensitive Sequence Comparison), LALIGN (Local Sequence Alignment), LCP (Local Content Program), MACAW (Multiple Alignment Construction & Analysis Workbench), MAP (Multiple Alignment Program), MBLKP, MBLKN, PIMA (Pattern-Induced Multi-sequence Alignment), SAGA (Sequence Alignment by Genetic Algorithm) and WHAT-IF. Such alignment programs can also be used to screen genome databases to identify polynucleotide sequences having substantially identical sequences. A number of genome databases are available, for example, a substantial portion of the human genome is available as part of the Human Genome Sequencing Project (J. Roach, http://weber.u.Washington.edu/~roach/human_genome_progress2.html weber.u.Washington.edu/~roach/human_genome_progress2.html) (Gibbs, 1995). At least twenty-one other genomes have already been sequenced, including, for example, *M. genitalium* (Fraser *et al.*, 1995), *M. jannaschii* (Bult *et al.*, 1996), *H. influenzae* (Fleischmann *et al.*, 1995), *E. coli* (Blattner *et al.*, 1997), and yeast (*S. cerevisiae*) (Mewes *et al.*, 1997), and *D. melanogaster* (Adams *et al.*, 2000). Significant progress has also been made in sequencing the genomes of model

organism, such as mouse, *C. elegans*, *Arabidopsis sp.* and *D. melanogaster*. Several databases containing genomic information annotated with some functional information are maintained by different organization, and are accessible via the internet, for example, ~~<http://www.tigr.org/tdb>~~; ~~<http://www.genetics.wisc.edu>~~; ~~<http://genome-www.stanford.edu/~ball>~~; ~~<http://hiv-web.lanl.gov>~~; ~~<http://www.ncbi.nlm.nih.gov>~~; ~~<http://www.ebi.ac.uk>~~; ~~<http://Pasteur.fr/other/biology>~~; and ~~<http://www.genome.wi.mit.edu>~~ www.tigr.org/tdb; www.genetics.wisc.edu; genome-www.stanford.edu/~ball; hiv-web.lanl.gov; www.ncbi.nlm.nih.gov; www.ebi.ac.uk; Pasteur.fr/other/biology; and www.genome.wi.mit.edu.